

LINEAR SYSTEMS IN $\mathbb{P}^3$ WITH LOW DEGREES AND LOW MULTIPLICITIES

MARCIN DUMNICKI

*Institute of Mathematics, Jagiellonian University,
ul. Grota-Roweckiego 26, 30-348 Kraków, Poland
Email address: Marcin.Dumnicki@im.uj.edu.pl*

ABSTRACT. We prove that the linear system of hypersurfaces in $\mathbb{P}^3$ of degree d , $14 \leq d \leq 40$, with double, triple and quadruple points in general position are non-special. This solves the cases that have not been completed in a paper by E. Ballico and M.C. Brambilla.
Keywords: linear systems, fat points.

1. INTRODUCTION

In what follows we assume that the ground field $\mathbb{K}$ is of characteristic zero. Let $d \in \mathbb{Z}$, let $m_1, \dots, m_r \in \mathbb{N}$. By $\mathcal{L}_3(d; m_1, \dots, m_r)$ we denote the linear system of hypersurfaces (in $\mathbb{P}^3 := \mathbb{P}^3(\mathbb{K})$) of degree d passing through r points $p_1, \dots, p_r$ in general position with multiplicities at least $m_1, \dots, m_r$ (the point with multiplicity m will often be called an m -point). The dimension of such system is denoted by $\dim \mathcal{L}_3(d; m_1, \dots, m_r)$. Define the *virtual dimension* of $L = \mathcal{L}_3(d; m_1, \dots, m_r)$

$$\text{vdim } L := \binom{d+3}{3} - \sum_{j=1}^r \binom{m_j+2}{3} - 1$$

and the *expected dimension* of L

$$\text{edim } L := \max\{\text{vdim } L, -1\}.$$

Observe that $\dim L \geq \text{edim } L$. If this inequality is strict then L is called *special*, *non-special* otherwise.

We will use the following notation: $m^{\times k}$ denotes the sequence of m 's taken k times,

$$m^{\times k} = \underbrace{(m, \dots, m)}_k.$$

2. RESULTS FOR MULTIPLICITIES BOUNDED BY 4

In [1] then following (see Thm. 1) was proven:

Theorem 1. *System $L = \mathcal{L}_3(d; 4^{\times x}, 3^{\times y}, 2^{\times z})$ is non-special for non-negative integers d, x, y, z satisfying $d \geq 41$.*

To consider all the lower cases, i.e. $d \leq 40$, authors of [1] proposed the algorithm, which essentially is the following:

Algorithm A.

```
-----
d=  -- degree should be chosen
N=binomial(d+3,3);
for z from 0 to ceiling(N/4) do
  for y from 0 to ceiling(N/10) do
```

```

for x from 0 to ceiling(N/20) do (
  if ((20*x+10*y+4*z>N-4)and(20*x+10*y+4*z<N+20)) then (
    -- computation of the rank of the interpolation matrix
    -- the rank is maximal if and only if the system is non-special
  )
)
)

```

With aid of the program Macaulay2 the following was proven ([1], Thm. 14):

Theorem 2. *System $L = \mathcal{L}_3(d; 4^{\times x}, 3^{\times y}, 2^{\times z})$ is non-special for non-negative integers d, x, y, z satisfying $9 \leq d \leq 13$.*

In order to prove Theorem 2 one has to check large number of cases, each case being computation of the rank of the interpolation matrix. The computation can be performed over $\mathbb{Z}_p$ for a small prime p , but still it is time-consuming. The case $d = 14$ (and all the next; for $d = 14$ we have 6816 nearly-square matrices of size 680) has not been done in [1] due to the length of computations. In this note we show how, using glueing theorem from [2], one can lower the number of cases. For example, one need to consider 261 cases for $d = 14$. For greater d 's the difference is even more visible — originally, for $d = 40$, 2294011 cases were needed, while in our approach it suffices to deal with 22 only.

The size of the each matrix is at most (about) 12341, which is small enough to be managed by a fast computer.

3. MAIN RESULT

Using a smarter version of Algorithm A we will show the following:

Theorem 3. *System $L = \mathcal{L}_3(d; 4^{\times x}, 3^{\times y}, 2^{\times z})$ is non-special for non-negative integers d, x, y, z satisfying $14 \leq d \leq 40$.*

We will use the following fact (see Thm. 1 and 9 in [2]):

Theorem 4. *Let $\mathcal{L}_3(k; \ell_1^{\times s_1}, \ell_2^{\times s_2})$ be non-special, let*

$$\begin{aligned}
 L_1 &= \mathcal{L}_3(d; m_1, \dots, m_r, \ell_1^{\times s_1}, \ell_2^{\times s_2}), \\
 L_2 &= \mathcal{L}_3(d; m_1, \dots, m_r, k+1).
 \end{aligned}$$

If either $-1 \leq \text{vdim } L_2 \leq \text{vdim } L_1$ or $\text{vdim } L_1 \leq \text{vdim } L_2 \leq -1$ then in order to show non-specialty of L_1 it is enough to show non-specialty of L_2 .

The above Theorem allows to “glue” s_1 ℓ_1 -points and s_2 ℓ_2 -points to one $(k+1)$ -point (the glueing will be denoted by $\ell_1^{\times s_1}, \ell_2^{\times s_2} \longrightarrow k+1$) during computations. We will glue according to the following

Proposition 5. *Systems $\mathcal{L}_3(3; 2^{\times 5})$ and $\mathcal{L}_3(9; 4^{\times a}, 3^{\times b})$ for non-negative integers a, b satisfying $2a + b = 22$ are non-special of virtual dimension equal to -1 .*

Proof. Non-specialty of $\mathcal{L}_3(3; 2^{\times 5})$ can be done by matrix computation or, without it, by [3]. Non-specialty of $\mathcal{L}_3(9; 4^{\times a}, 3^{\times b})$ follows from Thm. 2. The computation of vdim is straightforward:

$$\begin{aligned}
 \text{vdim } \mathcal{L}_3(9; 4^{\times a}, 3^{\times b}) &= \binom{13}{3} - a \binom{6}{3} - b \binom{5}{3} - 1 = 220 - 10(2a + b) - 1 = -1, \\
 \text{vdim } \mathcal{L}_3(3; 2^{\times 5}) &= \binom{6}{3} - 5 \binom{4}{3} - 1 = -1.
 \end{aligned}$$

□

So the possible glueings are $2^{\times 5} \rightarrow 4$ and $4^{\times a}, 3^{\times b} \rightarrow 10$ whenever $2a + b = 22$. Observe that the virtual dimension of a system before and after glueing does not change. Hence, it is enough to consider systems $\mathcal{L}_3(d; 10^{\times q}, 4^{\times x}, 3^{\times y}, 2^{\times z})$ with $2x + y \leq 21$ and $z \leq 4$. We summarize the above in the following algorithm:

Algorithm B.

```

-----
d=  -- degree should be chosen
N=binomial(d+3,3);
for z from 0 to 4 do
  for y from 0 to 10 do
    for x from 0 to 21 do
      (*) for q from 0 to ceiling(N/220) do (
        if ((220*q+20*x+10*y+4*z>N-4)and(220*q+20*x+10*y+4*z<N+20)and(2*a+b<22)) then (
          -- computation of the rank of the interpolation matrix
          -- the rank is maximal if and only if the system is non-special
        )
      )
    )
  )
-----

```

The above algorithm works properly for $d \geq 22$. For lower values of d the 10-points forces systems to be special and our approach does not work. To avoid this, we can change one line in Algorithm B

(*) for q from 1 to 1 do (-- 1 can be change into arbitrary non-negative integer choosing the number of 10-points to be considered. It is reasonable to choose this number as big as possible to obtain less cases. The right choices are one 10-point for $d \in \{13, \dots, 18\}$, five 10-points for $d = 19$, seven for $d = 20$ and eight for $d = 21$.

The above algorithm has been implemented in FreePascal (but it is easy to implement it in Singular or Macaulay2 or any other computer algebra program) and ran on several computers simultaneously. All systems appeared to be non-special.

Remark 6. By Thm. 1 (“splitting theorem”) from [2] we can avoid some of harmful computations. As an example consider $L = \mathcal{L}_3(40; 10^{\times 56}, 4^{\times 2})$. This system appears in a list of systems to be checked for $d = 40$. By splitting, it is enough to check non-specialty of $\mathcal{L}_3(39; 10^{\times 52}, 4^{\times 2})$ and $\mathcal{L}_3(40; 40, 10^{\times 4})$. The first system is assumed to be done during the previous stage (for $d = 39$), the second is non-special by [3]. Another possibility to make computations faster is to assume that four points are fundamental ones $((1 : 0 : 0 : 0), (0 : 1 : 0 : 0), (0 : 0 : 1 : 0)$ and $(0 : 0 : 0 : 1))$. This is possible by using a linear automorphism of $\mathbb{P}^3$. But then the size of the matrix is smaller by $\sum_{j=1}^4 \binom{m_j+2}{3}$, so, in most cases, where at least four 10-points are present, the advantage is 880 rows and columns. For d large enough we can also assume the multiplicities $m_1, \dots, m_4$ to be 14 (resp. 15, 18, 20), since the system $\mathcal{L}_3(13; 4^{\times a}, 3^{\times b})$ for $2a + b = 56$ (resp. $\mathcal{L}_3(14; 4^{\times a}, 3^{\times b})$ for $2a + b = 68$, $\mathcal{L}_3(17; 4^{\times a}, 3^{\times b})$ for $2a + b = 114$, $\mathcal{L}_3(19; 4^{\times a}, 3^{\times b})$ for $2a + b = 154$) is non-special of virtual dimension equal to -1 .

REFERENCES

- [1] E. Ballico, M.C. Brambilla, *Postulation of general quartuple fat point schemes in $\mathbb{P}^3$* , arXiv:math.AG/0810.1372v1 (2008), to appear in J. Pure App. Algebra
- [2] M. Dumnicki, *An algorithm to bound the regularity and non-emptiness of linear systems in $\mathbb{P}^n$* , arXiv:math.AG/0802.0925 (2008)
- [3] C. De Volder, A. Laface, *On linear systems of $\mathbb{P}^3$ through multiple points*, J. Algebra 310, 207–217 (2007)